



Lysophosphatidic Acid: Friend or Foe of the Ovarian Surface Epithelium

Darren J. Park PhD, Kendra D. Martyn PhD, and Bonnie J. Warn-Cramer PhD
Natural Products and Cancer Biology Program
Cancer Research Center of Hawaii

Introduction

Ovarian cancer accounts for more deaths among women than any other cancer of the female reproductive system, and ranks fourth in cancer-causing deaths among women. The American Cancer Society estimates that in 2005, approximately 22,220 new cases of ovarian cancer will be diagnosed, and there will be about 16,210 deaths from ovarian cancer in the United States¹. Over ninety percent of human ovarian adenocarcinomas arise from the single layer of ovarian surface epithelial (OSE) cells that surround the ovaries². It is thought that the cyclic rounds of rupture and repair of the ovarian surface epithelium induced by ovulation, predispose these cells to neoplasia^{3,4}.

Lysophosphatidic acid (LPA) is a phospholipid growth factor found in serum, follicular and ascites fluid. LPA plays an integral part in repairing the wound caused by follicular rupture of the epithelial surface during ovulation by stimulating OSE cell division, resulting in rapid closure and repair of the wound. LPA initiates a

signaling pathway within the cells of the ovarian surface epithelium that induces enhanced cell proliferation (Figure 1). This signaling results in the down-regulation of the growth and tumor suppressor functions of a channel-forming protein, connexin43 (Cx43), which then permits an increase in cell proliferation.

These cycles of LPA-mediated down-regulation of Cx43 function, which assist in quick repair of the ovarian surface after ovulation, may also allow abnormal, pre-cancerous OSE cells to proliferate and form tumors by escaping the tumor suppressing properties of Cx43. LPA also plays an important role in promoting the vascular network required for the growth of ovarian tumors, and it is known to increase resistance of the tumor cells to chemotherapeutic agents⁵. Clinical trials are currently ongoing to determine the effectiveness of measuring the plasma levels of LPA in the detection of ovarian cancer since ovarian cancer cells and not normal OSE cells secrete LPA, resulting in elevated plasma levels of LPA in about 90% of women with early stages ovarian cancer⁶.




Figure 1.— LPA signaling in normal rat ovarian surface epithelial (ROSE) cells.

LPA initiates signaling by binding to an extracellular site on the Edg2 transmembrane-receptor, leading to the G_i-mediated transactivation of the epidermal growth factor receptor (EGFR), also a transmembrane receptor. Transactivation of the EGFR leads to the activation of Erk kinases, increased phosphorylation of Cx43, and the down-regulation of GJC. The down-regulation of Cx43's growth-suppressive function allows enhanced cell proliferation that aids in wound repair. However, the enhanced cell proliferation may also allow abnormal cells to escape from the normal tumor-suppressor function of Cx43 and thus proliferate and contribute to the development of ovarian cancer.

Connexin Expression In the Ovarian Surface Epithelial Cells

Connexins (Cxs) mediate gap junctional communication (GJC) by forming aqueous channels between neighboring cells and facilitating the passive exchange of small molecules and ions, less than one kilodalton in mass. Cxs are integral plasma membrane proteins with cytoplasmically localized N- and C- terminal domains that create a single intracellular loop. To form a complete intercellular channel, each cell contributes a Cx hexamer, or a half channel, that docks with a hexamer from a neighboring cell.

Since the reduction or loss of GJC and/or Cx expression is associated with enhanced cell proliferation in tumor cells as compared to normal cells^{7,8}, Cxs are thought to have a tumor suppressor function. Likewise, inducing the expression of Cx43 in tumor cells and in Cx43-null cells decreases rates of cell proliferation⁹. Importantly, forced expression of Cx43 in Cx43-deficient ovarian carcinoma cells increases the cells' dependence on serum growth factors and reduces their tumorigenicity in nude mice¹⁰. Normal rat OSE (ROSE) cells and normal human OSE (HOSE) cells primarily express the Cx43 protein, whereas Cx43 and GJC are greatly reduced or absent in neoplastic derivatives of ROSE cells and in human ovarian adenocarcinoma cells¹¹. These data support the role of the Cx43 gap junction protein, in tumor suppression of OSE cells. However, the precise mechanisms by which Cx43 exerts its anti-proliferative effect are not yet understood.

LPA-Mediated Signaling In Ovarian Surface Epithelial Cells

LPA affects the regulation of cell growth, adhesion, and migration by engaging the Edg2/LPA₁, Edg4/LPA₂, and Edg7/LPA₃ transmembrane receptors present on the surface of normal and transformed ovarian cells¹². The Edg receptors couple to G proteins (GPCRs, G protein-coupled receptors) comprised of alpha, beta, and gamma subunits that initiate unique signaling pathways. The activation of a particular signaling pathway is determined by the different G-alpha subclasses, either G_s, G_i, G_q, or G_{12/13}. The Edg2 receptor primarily signals through a G_i-dependent pathway and the Edg7 receptor through a G_q-dependent pathway while the Edg4 receptor signals through G_i- and G_q-coupled pathways in OSE cells¹³.

In some cell types, the transactivation of the epidermal growth factor receptor (EGFR)/Her-1 is required for mediating the LPA-induced activation of extracellular-signal-regulated kinases (Erks)¹². The EGFR gene is often amplified, over expressed, or mutated in ovarian carcinomas, resulting in various alterations of the EGFR signaling pathways, including ligand-independent EGFR signaling¹³. Historically, poor prognosis has been associated with over expression of the EGFR in ovarian carcinomas, however recent univariate and multivariate analysis has demonstrated a lack of correlation between EGFR over expression and prognosis¹⁴⁻¹⁶. Alterations in EGFR-mediated signaling pathways have been implicated in human cancer,¹⁷ and antibody therapies directed against the EGFR are the subject of ongoing clinical research¹⁸. These trials show promise in the treatment of ovarian cancer.

LPA Transiently Down-Regulates Connexin43 Function in Ovarian Surface Epithelial Cells

The EGFR is known to activate Erk kinases, which have been shown

to play a key role in the disruption of GJC and the phosphorylation of Cx43¹⁹⁻²¹. However, Postma et al. have suggested that the non-receptor tyrosine kinase, c-Src, and not Erk kinases, are critical for the disruption of GJC induced by LPA in rat fibroblast cells; however, tyrosine phosphorylation of Cx43 was not detected and the precise mechanism for the c-Src-mediated disruption of GJC was not determined²². We have demonstrated in normal rat OSE cells that LPA-induced transactivation of the EGFR is required for the transient down-regulation of Cx43-mediated GJC, the activation of Erk5, and the maximal increase in Cx43 phosphorylation (Lin et al., manuscript in revision). These results may support the conclusions of Cameron et al., which suggest that Erk5 is more important than the more studied Erk1 and Erk2 kinases²¹ in cells treated with epidermal growth factor.

Conclusions

To better understand the pathogenesis and development of ovarian cancer, studies of the LPA-mediated signaling pathways that are activated in wound repair in response to follicular rupture in normal OSE cells are required. It is thought that Cx-mediated GJC between neighboring cells is important for coordinating the responses of different cell types in wound repair and it has been shown that the down-regulation of Cx43 increases the rate of wound closure and reduces inflammation and scarring in a skin wound-repair model²³. Although the transient down-regulation of Cx43 by LPA may play an important role in enhancing normal OSE cell proliferation for repair of the ovarian epithelium after follicular rupture, the down-regulation of GJC by LPA may also allow aberrant cells that naturally arise during cell proliferation to escape from the normal growth-suppressive effects of Cx43. This possible course of events could contribute to the development of ovarian cancer. Considering these two possible outcomes resulting from LPA's down-regulation of Cx43 function in OSE cells, LPA may act either as a friend or a foe of the ovarian surface epithelium.

For more information on the Cancer Research Center of Hawaii, please visit our website at www.crch.org.

Acknowledgements

Supported by an American Cancer Society Institutional Research Grant (IRG-92-025-08-IRG to BWC) and grants from the NIH (RR-16453, Charles Boyd, P.I.; BWC, project P.I. and P20-RR16467, HS-BRIN for travel awards to BWC and DJP).

References

1. American Cancer Society Cancer Facts and Figures 2005.
2. Murdoch WJ. Ovarian surface epithelium, ovulation and carcinogenesis. *Biol Rev Camb Philos Soc*. Nov 1996;71(4):529-543.
3. Testa JR, Gettis LA, Salazar H, et al. Spontaneous transformation of rat ovarian surface epithelial cells results in well to poorly differentiated tumors with a parallel range of cytogenetic complexity. *Cancer Res*. May 15 1994;54(10):2778-2784.
4. Fathalla MF. Incessant ovulation--a factor in ovarian neoplasia? *Lancet*. Jul 17 1971;2(7716):163.
5. Frankel A, Mills GB. Peptide and lipid growth factors decrease cis-diamminedichloroplatinum-induced cell death in human ovarian cancer cells. *Clin Cancer Res*. Aug 1996;2(8):1307-1313.
6. Sutphen R, Xu Y, Wilbanks GD, et al. Lysophospholipids are potential biomarkers of ovarian cancer. *Cancer Epidemiol Biomarkers Prev*. Jul 2004;13(7):1185-1191.
7. Yamasaki H, Naus CC. Role of connexin genes in growth control. *Carcinogenesis*. Jun 1996;17(6):1199-1213.
8. Trosko JE, Ruch RJ. Cell-cell communication in carcinogenesis. *Front Biosci*. Feb 15 1998;3:d208-236.
9. Martyn KD, Kurata WE, Warn-Cramer BJ, Burt JM, TenBroek E, Lau AF. Immortalized connexin43 knockout cell lines display a subset of biological properties associated with the transformed phenotype. *Cell Growth Differ*. Sep 1997;8(9):1015-1027.

See *Lysophosphatidic Acid*, p. 169

10. Fernstrom MJ, Koffler LD, Abou-Rjaily G, Boucher PD, Shewach DS, Ruch RJ. Neoplastic reversal of human ovarian carcinoma cells transfected with connexin43. *Exp Mol Pathol*. Aug 2002;73(1):54-60.
11. Hanna EA, Umhauer S, Roshong SL, et al. Gap junctional intercellular communication and connexin43 expression in human ovarian surface epithelial cells and ovarian carcinomas in vivo and in vitro. *Carcinogenesis*. Jul 1999;20(7):1369-1373.
12. Goetzl EJ, Dolezalova H, Kong Y, et al. Distinctive expression and functions of the type 4 endothelial differentiation gene-encoded G protein-coupled receptor for lysophosphatidic acid in ovarian cancer. *Cancer Res*. Oct 15
13. Anliker B, Chun J. Lysophospholipid G protein-coupled receptors. *J Biol Chem*. May 14 2004;279(20):20555-20558.
14. Rubio I, Rennert K, Wittig U, Wetzker R. Ras activation in response to lysophosphatidic acid requires a permissive input from the epidermal growth factor receptor. *Biochem J*. Dec 15 2003;376(Pt 3):571-576.
15. Abdollahi A, Gruver BN, Patriotis C, Hamilton TC. Identification of epidermal growth factor-responsive genes in normal rat ovarian surface epithelial cells. *Biochem Biophys Res Commun*. Jul 18 2003;307(1):188-197.
16. Nielsen JS, Jakobsen E, Holund B, Bertelsen K, Jakobsen A. Prognostic significance of p53, Her-2, and EGFR overexpression in borderline and epithelial ovarian cancer. *Int J Gynecol Cancer*. Nov-Dec 2004;14(6):1086-1096.
17. Lee CH, Huntsman DG, Cheang MC, et al. Assessment of Her-1, Her-2, and Her-3 expression and Her-2 amplification in advanced stage ovarian carcinoma. *Int J Gynecol Pathol*. Apr 2005;24(2):147-152.
18. Elie C, Geay JF, Morcos M, et al. Lack of relationship between EGFR-1 immunohistochemical expression and prognosis in a multicentre clinical trial of 93 patients with advanced primary ovarian epithelial cancer (GINECO group). *Br J Cancer*. Aug 2 2004;91(3):470-475.
19. Downward J, Yarden Y, Mayes E, et al. Close similarity of epidermal growth factor receptor and v-erb-B oncogene protein sequences. *Nature*. Feb 9-15 1984;307(5951):521-527.
20. Nicodemus CF, Berek JS. Monoclonal antibody therapy of ovarian cancer. *Expert Rev Anticancer Ther*. Feb 2005;5(1):87-96.
21. Warn-Cramer BJ, Cottrell GT, Burt JM, Lau AF. Regulation of connexin-43 gap junctional intercellular communication by mitogen-activated protein kinase. *J Biol Chem*. Apr 10 1998;273(15):9188-9196.
22. Hossain MZ, Ao P, Boynton AL. Rapid disruption of gap junctional communication and phosphorylation of connexin43 by platelet-derived growth factor in T51B rat liver epithelial cells expressing platelet-derived growth factor receptor. *J Cell Physiol*. Jan 1998;174(1):66-77.
23. Cameron SJ, Malik S, Akaike M, et al. Regulation of epidermal growth factor-induced connexin 43 gap junction communication by big mitogen-activated protein kinase1/ERK5 but not ERK1/2 kinase activation. *J Biol Chem*. May 16 2003;278(20):18682-18688.
24. Postma FR, Hengeveld T, Alblas J, et al. Acute loss of cell-cell communication caused by G protein-coupled receptors: a critical role for c-Src. *J Cell Biol*. Mar 9 1998;140(5):1199-1209.
25. Qiu C, Coutinho P, Frank S, et al. Targeting connexin43 expression accelerates the rate of wound repair. *Curr Biol*. Sep 30 2003;13(19):1697-1703.

Classified Notices

To place a classified notice:

HMA members.—As a benefit of membership, HMA members may place a complimentary one-time classified ad in HMJ as space is available.

Nonmembers.—Rates are \$1.50 a word with a minimum of 20 words or \$30. Not commissionable.

For more information call (808) 536-7702.

Office Space & Support Services

ALA MOANA BLDG.—PHYSICIAN WANTED to share space and support services. Interest in physical rehab. preferred. We have unique time-share arrangements starting at one half-day per week. Run your practice with no fixed overhead. Contact Dr. Speers, REHABILITATION ASSOCIATES, 955-7244.

Physician Wanted

NEPHROLOGIST—REQUIRES MD, completion of fellowship in Nephrology, completion of residency in Internal Med., BE/BC Nephrology, BC Internal Med., eligibility for HI med. license. Location: Kahului, HI. Mail CV to Maui Nephrology, P.O. Box 3260, Wailuku, HI 96793.

CT Consulting

STARTING A NEW PRACTICE? —CT CONSULTING, INC., provides business insight to effectively contract with healthplans, establish billing guidelines, interview staff and coordinate other business functions. With 15+ years of experience in the healthcare industry, CT Consulting adds business savvy to your practice. Contact Catherine at (808) 330-7738. e-mail: ctoconsulting@hawaii.rr.com.

Surfer's Medical Association: Conference in Biarritz, France

The first European meeting of the SMA will be held from September 30 to October 9, 2005. The conference would be of interest to physicians and allied health-care professionals who treat surfers as patients and/or who are surfers themselves.

The conference program will include presentations by all attendees to update the status of the health-care/surfing interface plus networking with international colleagues. Additional details are available on the SMA website (www.damoon.net) in the events section.

The conference registration fee of \$2,200 USD includes one oceanfront room (based on double occupancy) for ten nights, three meals per day, and ground transport each day to selected surf spots.

To enroll, make check payable to SMA:BIARRITZ and mail to 1330 Ala Moana Blvd., #2101, Honolulu, HI 96814. Questions may be directed to the conference chairperson, Dr. Bob Speers at speers@lava.net.